

Oregon 2002 -2005 BRFSS Data - Behavior Risk Survey Topics by County

Data are adjusted to the 2000 U.S. Census Population.	
County	% who are obese (body mass index >30 kg/m2)
OREGON	22.1%
Baker	18.0%
Benton	16.4%
Clackamas	21.1%
Clatsop	25.7%
Columbia	30.9%**
Coos	23.1%
Crook	23.8%
Curry	15.2%**
Deschutes	18.3%**
Douglas	27.0%**
Grant	22.9%
Harney	26.2%
Hood River	19.8%
Jackson	20.9%
Jefferson	28.7%
Josephine	23.3%
Klamath	23.2%
Lake	24.1%
Lane	23.0%
Lincoln	28.0%**
Linn	30.5%**
Malheur	30.1%**
Marion	25.1%**
Morrow	29.3%
Multnomah	19.9%**
Polk	21.0%
Tillamook	20.7%
Umatilla	26.1%
Union	20.7%
Wallowa	9.9%**
Washington	20.4%
Yamhill	25.9%
Gilliam/Wheeler	43.1%†
Sherman/Wasco	23.4%

Age-adjusted Prevalence of Selected Chronic Diseases and Related Risk Factors Among Adults by County, Oregon 2002-2005

Tables were created using combined BRFSS survey results from 2002 through 2005. Two numbers are provided for each condition or disease: an unadjusted rate and an age-adjusted rate.

Unadjusted rates, provided in separate tables, give a description of the absolute burden of a disease or risk factor for an individual county.

Age-adjusted rates supplied in this table allow you to compare rates for a county to the State's rate. They adjust for differences that could result from a county's population being, on average, older or younger than the State's. When comparing counties with Oregon or with each other, use these age-adjusted rates.

† Rate based on less than 50 respondents overall or fewer than 12 persons in one age group; may not accurately reflect behavior of entire county and results should be interpreted with caution. Estimates based on 10 or fewer respondents are not reported.

** Statistically significant difference compared to Oregon.

Three age groups were used for age adjusting: 18-34yrs., 35-54yrs, and 55+yrs.

Source: Oregon Behavioral Risk Factor Surveillance System - healthoregon.org/brfss