

Oregon 2002 -2005 BRFSS Data - Behavior Risk Survey Topics by County

Data are adjusted to the 2000 U.S. Census Population.	
County	% who had a mammogram within past 2 years (women 40+ years old)
OREGON	73.3%
Baker	74.2%†
Benton	80.0%
Clackamas	75.6%
Clatsop	65.5%
Columbia	72.6%
Coos	72.2%
Crook	73.1%†
Curry	84.5%†
Deschutes	68.2%
Douglas	68.2%
Grant	66.8%†
Harney	71.1%†
Hood River	84.3%
Jackson	69.7%
Jefferson	61.8%†
Josephine	61.2%**
Klamath	81.9%
Lake	81.7%†
Lane	74.8%
Lincoln	63.2%
Linn	64.1%
Malheur	63.6%†
Marion	76.6%
Morrow	88.4%†
Multnomah	75.8%
Polk	74.1%
Tillamook	54.7%**
Umatilla	70.3%
Union	75.1%
Wallowa	60.6%†
Washington	75.2%
Yamhill	72.2%
Gilliam/Wheeler	--
Sherman/Wasco	77.1%

Age-adjusted Prevalence of Selected Chronic Diseases and Related Risk Factors Among Adults by County, Oregon 2002-2005

Tables were created using combined BRFSS survey results from 2002 through 2005. Two numbers are provided for each condition or disease: an unadjusted rate and an age-adjusted rate.

Unadjusted rates, provided in separate tables, give a description of the absolute burden of a disease or risk factor for an individual county.

Age-adjusted rates supplied in this table allow you to compare rates for a county to the State's rate. They adjust for differences that could result from a county's population being, on average, older or younger than the State's. When comparing counties with Oregon or with each other, use these age-adjusted rates.

† Rate based on less than 50 respondents overall or fewer than 12 persons in one age group; may not accurately reflect behavior of entire county and results should be interpreted with caution. Estimates based on 10 or fewer respondents are not reported.

** Statistically significant difference compared to Oregon.

Three age groups were used for age adjusting: 18-34yrs., 35-54yrs, and 55+yrs.