

Oregon 2002 -2005 BRFSS Data - Behavior Risk Survey Topics by County

Data are adjusted to the 2000 U.S. Census Population.	
County	% who met CDC recommendations for physical activity
OREGON	54.7%
Baker	50.8%
Benton	58.2%
Clackamas	54.6%
Clatsop	57.3%
Columbia	56.5%
Coos	60.4%
Crook	55.5%
Curry	67.9%**†
Deschutes	57.5%
Douglas	52.1%
Grant	72.9%†
Harney	56.0%†
Hood River	44.7%†
Jackson	58.0%
Jefferson	53.9%
Josephine	56.1%
Klamath	54.1%
Lake	63.3%†
Lane	58.8%
Lincoln	50.1%
Linn	54.9%
Malheur	47.1%
Marion	49.6%**
Morrow	40.4%
Multnomah	56.0%
Polk	57.9%
Tillamook	52.5%
Umatilla	37.4%**
Union	61.8%
Wallowa	51.8%†
Washington	51.3%
Yamhill	55.5%
Gilliam/Wheeler	64.9%†
Sherman/Wasco	64.5%

Age-adjusted Prevalence of Selected Chronic Diseases and Related Risk Factors Among Adults by County, Oregon 2002-2005

Tables were created using combined BRFSS survey results from 2002 through 2005. Two numbers are provided for each condition or disease: an unadjusted rate and an age-adjusted rate.

Unadjusted rates, provided in separate tables, give a description of the absolute burden of a disease or risk factor for an individual county.

Age-adjusted rates supplied in this table allow you to compare rates for a county to the State's rate. They adjust for differences that could result from a county's population being, on average, older or younger than the State's. When comparing counties with Oregon or with each other, use these age-adjusted rates.

† Rate based on less than 50 respondents overall or fewer than 12 persons in one age group; may not accurately reflect behavior of entire county and results should be interpreted with caution. Estimates based on 10 or fewer respondents are not reported.

** Statistically significant difference compared to Oregon.

Three age groups were used for age adjusting: 18-34yrs., 35-54yrs, and 55+yrs.