

Table III: Age-Adjusted and Unadjusted Prevalence of Preventive Health Screening among Adults, by County, Oregon 2006-2009

County	% who had their cholesterol checked within past five years (≥ 18 years old)		% who had a PAP test within past three years (women 18-65 years old with intact cervix)	
	unadjusted	age-adjusted	unadjusted	age-adjusted
OREGON		71.3%		85.8%
Baker	72.8%	66.8%	76.0%	75.8%
Benton	70.2%	75.1%	97.0%	95.1%*
Clackamas	72.4%	70.6%	88.7%	88.3%
Clatsop	72.7%	68.4%	56.4%	61.3%*
Columbia	72.3%	67.8%	89.6%	89.8%
Coos	80.9%	75.1%	78.5%	78.3%
Crook	83.0%	81.7%	–	–
Curry	65.6%	49.7%*	–	–
Deschutes	76.2%	73.3%	80.3%	79.8%
Douglas	77.8%	73.6%	80.1%	79.9%
Grant	76.7%	54.2%*	–	–
Harney	67.3%	58.6%	–	–
Hood River	69.8%	69.0%	94.8%	92.8%
Jackson	73.9%	71.1%	82.2%	85.5%
Jefferson	59.0%	56.3%*	64.8%	63.1%*
Josephine	79.7%	75.3%	81.7%	81.6%
Klamath	74.4%	72.2%	81.2%	81.1%
Lake	74.5%	69.6%	–	–
Lane	68.3%	67.3%	85.2%	86.1%
Lincoln	65.5%	54.1%*	82.9%	85.1%
Linn	70.6%	68.1%	83.3%	83.3%
Malheur	76.4%	74.8%	89.8%	88.4%
Marion	73.1%	73.0%	82.7%	84.3%
Morrow	52.9%	44.5%*	–	–
Multnomah	71.8%	71.6%	86.6%	86.6%
Polk	73.4%	72.1%	77.1%	77.6%
Tillamook	76.1%	70.1%	86.8%	87.9%
Umatilla	73.6%	73.3%	82.6%	80.3%
Union	71.8%	70.2%	89.8%	89.9%
Wallowa	82.5%	65.2%*	–	–
Washington	73.7%	74.0%	91.5%	91.5%
Wheeler	–	–	–	–
Yamhill	70.7%	71.4%	83.9%	83.2%
Gilliam/Sherman/Wasco	70.3%	66.5%	–	–

* Statistically significant difference compared with Oregon.

† This number may be statistically unreliable and should be interpreted with caution.

– This number is suppressed because it is statistically unreliable.

Age-adjusted estimates are adjusted to the 2000 Standard Population using three age groups (18-34, 35-54, and 55+).

Source: Oregon BRFSS County Combined Dataset 2006-2009

Table III (continued): Age-Adjusted and Unadjusted Prevalence of Preventive Health Screening among Adults, by County, Oregon 2006-2009

County	% who had a mammogram within past two years (women 50-74 years old)	% who had an FOBT [^] in the past year or a Colonoscopy/ Sigmoidoscopy within the past five years (50-75 year olds)
OREGON	82.0%	56.8%
Baker	70.6%	47.0%
Benton	89.5%	68.1%
Clackamas	86.4%	57.3%
Clatsop	76.3%	44.1%
Columbia	84.3%	62.4%
Coos	89.6%	62.3%
Crook	–	47.5%
Curry	65.9%	61.9%
Deschutes	72.8%	59.0%
Douglas	77.3%	60.3%
Grant	–	58.6%
Harney	–	35.5%†
Hood River	–	26.3%†
Jackson	82.4%	54.6%
Jefferson	–	56.0%
Josephine	72.2%	48.6%
Klamath	75.2%	47.5%
Lake	–	–
Lane	82.8%	60.7%
Lincoln	63.8%*	46.8%
Linn	77.4%	57.9%
Malheur	70.5%	42.8%
Marion	85.9%	55.5%
Morrow	–	–
Multnomah	85.7%	57.9%
Polk	89.0%	60.0%
Tillamook	72.2%	46.2%
Umatilla	81.0%	53.7%
Union	71.8%	58.6%
Wallowa	–	56.3%
Washington	86.5%	58.3%
Wheeler	–	–
Yamhill	76.3%	59.8%
Gilliam/Sherman/Wasco	80.1%	57.7%

* Statistically significant difference compared with Oregon.

† This number may be statistically unreliable and should be interpreted with caution.

– This number is suppressed because it is statistically unreliable.

[^] Fecal Occult Blood Test

Source: Oregon BRFSS County Combined Dataset 2006-2009