

Table III: Age-adjusted and unadjusted prevalence of preventive health screening among adults, by county, Oregon 2008-2011

	Cholesterol checked within past five years (≥ 18 years old)		PAP test within past three years (women 21-65 years old with a cervix)	
	unadjusted	age-adjusted	unadjusted	age-adjusted
OREGON		73.1%		84.4%
Baker	74.2%	68.2%	76.3%	75.9%
Benton	74.4%	76.8%	92.1%	91.4%*
Clackamas	76.5%	74.5%	88.2%	87.2%
Clatsop	73.1%	67.8%	69.3%	71.3%
Columbia	74.2%	70.4%	77.6%	77.9%
Coos	80.1%	73.5%	73.3%	74.9%*
Crook	80.3%	78.3%	57.9%	60.1%
Curry	70.8%	58.1%	–	–
Deschutes	76.2%	73.4%	88.4%	87.7%
Douglas	72.4%	66.3%*	87.0%	87.3%
Grant	85.0%	67.6%	–	–
Harney	76.5%	74.1%	–	–
Hood River	77.4%	76.0%	91.9%	92.7%*
Jackson	73.2%	69.8%	81.9%	82.7%
Jefferson	73.3%	60.7%*	–	–
Josephine	79.0%	70.3%	84.7%	85.6%
Klamath	75.7%	73.3%	84.3%	83.7%
Lake	64.0%	56.5%	–	–
Lane	71.7%	69.8%	79.4%	79.9%
Lincoln	74.3%	66.3%	73.5%	73.6%
Linn	71.0%	68.8%	82.1%	81.6%
Malheur	74.4%	70.1%	74.3%	70.4%
Marion	73.9%	73.6%	83.8%	83.9%
Morrow	69.4%	65.9%	92.2%	93.9%*
Multnomah	76.1%	76.0%*	85.9%	86.2%
North Central HD	77.1%	73.2%	89.8%	90.4%
Polk	75.7%	74.7%	91.6%	90.5%
Tillamook	72.3%	63.5%	90.5%	89.7%
Umatilla	73.9%	72.3%	78.0%	76.3%
Union	85.8%	85.2%*	92.9%	91.3%
Wallowa	85.2%	65.4%*	–	–
Washington	75.1%	75.1%	87.6%	87.2%
Wheeler	–	–	–	–
Yamhill	71.2%	70.8%	82.5%	80.3%

* Statistically significant difference compared with all other counties (p-value ≤ 0.05)

† This number may be statistically unreliable and should be interpreted with caution.

– This number is suppressed because it is statistically unreliable.

Age-adjusted estimates are adjusted to the 2000 Standard Population using three age groups (18-34, 35-54, and 55+).

Source: Oregon BRFSS County Combined Dataset 2008-2011

Table III (continued): Age-adjusted and unadjusted prevalence of preventive health screening among adults, by county, Oregon 2008-2011

	Mammogram within past two years (women 50-74 years old)	Current [^] on colorectal cancer screening (50-75 year olds)
OREGON	79.7%	61.2%
Baker	71.8%	46.9%*
Benton	85.8%	69.3%*
Clackamas	82.5%	64.4%
Clatsop	78.0%	46.0%*
Columbia	77.8%	63.8%
Coos	76.2%	65.7%
Crook	58.8%	35.5%*
Curry	58.5%	57.7%
Deschutes	77.7%	65.0%
Douglas	75.8%	68.5%*
Grant	–	33.7%†
Harney	–	49.4%
Hood River	95.0%*	55.2%
Jackson	76.8%	57.8%
Jefferson	80.1%	51.2%
Josephine	69.0%*	53.1%*
Klamath	75.1%	58.0%
Lake	–	44.5%
Lane	82.4%	62.4%
Lincoln	68.7%*	57.8%
Linn	75.3%	61.2%
Malheur	56.6%*	64.0%
Marion	83.5%	61.2%
Morrow	–	40.5%
Multnomah	84.6%*	62.3%
North Central HD	68.9%	56.5%
Polk	86.9%	66.0%
Tillamook	74.5%	57.3%
Umatilla	77.6%	54.0%
Union	83.7%	41.3%*
Wallowa	–	70.4%
Washington	81.2%	64.7%
Wheeler	–	–
Yamhill	68.9%*	50.9%*

* Statistically significant difference compared with all other counties (p-value <= 0.05)

† This number may be statistically unreliable and should be interpreted with caution.

– This number is suppressed because it is statistically unreliable.

[^] Fecal occult blood test in the past year or a colonoscopy within 10 years or sigmoidoscopy within the past five years
Age-adjusted estimates are adjusted to the 2000 Standard Population using three age groups (18-34, 35-54, and 55+).

Source: Oregon BRFSS County Combined Dataset 2008-2011