

**TABLE 3. Estimated vaccination coverage among children aged 19-35 months, by selected individual vaccines and vaccination series\*, by state and local area – National Immunization Survey, United States, 2013<sup>†</sup>**

	MMR (≥1 doses)		DTaP (≥4 doses)		Hep B (birth) <sup>§</sup>		HepA (≥2 doses) <sup>¶</sup>		Rotavirus**		Combined Vaccine series*	
	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
<b>US National</b>	91.9	(±0.9)	83.1	(±1.3)	74.2	(±1.4) <sup>††</sup>	54.7	(±1.6)	72.6	(±1.5) <sup>††</sup>	70.4	(±1.5)
<b>HHS REGION X</b>	91.9	(±2.5) <sup>††</sup>	81.2	(±4.1)	71.6	(±4.3)	56.2	(±5.0)	72.1	(±4.4)	69.2	(±4.6)
<b>Oregon</b>	<b>89.4</b>	<b>(±4.4)</b>	<b>83.8</b>	<b>(±5.2)</b>	<b>66.8</b>	<b>(±6.3)</b>	<b>55.9</b>	<b>(±6.7)</b>	<b>64.3</b>	<b>(±6.7)</b>	<b>66.6</b>	<b>(±6.5)</b>

**Abbreviations:** CI = confidence interval; DTaP = diphtheria, tetanus toxoids and pertussis vaccines, diphtheria and tetanus toxoids, and diphtheria, tetanus toxoids and acellular pertussis vaccine; HepB = hepatitis B vaccine; Hib = *Haemophilus influenzae* type b vaccine; MMR = measles, mumps, and rubella vaccine; PCV = pneumococcal conjugate vaccine

\*Includes ≥4 doses of DTP/DT/DTaP, ≥3 doses of poliovirus vaccine, ≥1 doses of measles-containing vaccine, full series of Hib vaccine (3 or 4 doses, depending on product type), ≥3 doses of hepatitis B vaccine, ≥1 dose of varicella vaccine, and ≥4 doses of PCV.

<sup>†</sup>Children in the 2013 National Immunization Survey were born during January 2010-May 2012.

<sup>§</sup> 1 or more doses of hepatitis B vaccine administered between birth and age 3 days.

<sup>¶</sup> ≥2 doses hepatitis A vaccine and measured among children aged 19-35 months.

\*\*≥2 or ≥3 doses of rotavirus vaccine, depending on product type received (≥2 doses for Rotarix [RV1] and ≥3 doses for RotaTeq [RV5]).

<sup>††</sup> Statistically significant increase in coverage compared to 2012 (p<0.05).