

Oregon 2002 -2005 BRFSS Data - Behavior Risk Survey Topics by County

Data are adjusted to the 2000 U.S. Census Population.	
County	% who are overweight (body mass index 2529.9 kg/m2)
OREGON	37.0%
Baker	37.4%
Benton	37.5%
Clackamas	38.1%
Clatsop	33.9%
Columbia	35.6%
Coos	37.3%
Crook	44.8%
Curry	39.1%
Deschutes	38.8%
Douglas	36.9%
Grant	42.4%
Harney	49.4%**
Hood River	39.2%
Jackson	36.2%
Jefferson	42.7%
Josephine	33.2%
Klamath	39.2%
Lake	37.3%
Lane	35.7%
Lincoln	37.8%
Linn	35.0%
Malheur	40.1%
Marion	39.4%
Morrow	46.2%
Multnomah	34.8%**
Polk	37.0%
Tillamook	38.8%
Umatilla	40.4%
Union	40.0%
Wallowa	37.1%
Washington	37.1%
Yamhill	36.8%
Gilliam/Wheeler	24.0%†
Sherman/Wasco	34.9%

Age-adjusted Prevalence of Selected Chronic Diseases and Related Risk Factors Among Adults by County, Oregon 2002-2005

Tables were created using combined BRFSS survey results from 2002 through 2005. Two numbers are provided for each condition or disease: an unadjusted rate and an age-adjusted rate.

Unadjusted rates, provided in separate tables, give a description of the absolute burden of a disease or risk factor for an individual county.

Age-adjusted rates supplied in this table allow you to compare rates for a county to the State's rate. They adjust for differences that could result from a county's population being, on average, older or younger than the State's. When comparing counties with Oregon or with each other, use these age-adjusted rates.

† Rate based on less than 50 respondents overall or fewer than 12 persons in one age group; may not accurately reflect behavior of entire county and results should be interpreted with caution. Estimates based on 10 or fewer respondents are not reported.

** Statistically significant difference compared to Oregon.

Three age groups were used for age adjusting: 18-34yrs., 35-54yrs, and 55+yrs.