

**TABLE 7-9. Late Neonatal Deaths<sup>1</sup> by Weeks of Gestation and Weight  
Oregon Residents, Birth Cohort 2009**

Birthweight (In Grams)	Total	Weeks of Gestation									
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+	N.S.
<b>Total</b> .....	27	-	-	5	6	4	1	8	3	-	-
001-349 .....	-	-	-	-	-	-	-	-	-	-	-
350-499 .....	1	-	-	1	-	-	-	-	-	-	-
<500 .....	1	-	-	1	-	-	-	-	-	-	-
500-749 .....	2	-	-	2	-	-	-	-	-	-	-
750-999 .....	2	-	-	2	-	-	-	-	-	-	-
1000-1249 .....	5	-	-	-	4	1	-	-	-	-	-
1250-1499 .....	1	-	-	-	1	-	-	-	-	-	-
1500-1999 .....	2	-	-	-	1	1	-	-	-	-	-
2000-2499 .....	2	-	-	-	-	-	1	1	-	-	-
<2500 .....	15	-	-	5	6	2	1	1	-	-	-
2500+ .....	12	-	-	-	-	2	-	7	3	-	-
2500-2999 .....	3	-	-	-	-	2	-	1	-	-	-
3000-3499 .....	6	-	-	-	-	-	-	4	2	-	-
3500-3999 .....	3	-	-	-	-	-	-	2	1	-	-
4000-4499 .....	-	-	-	-	-	-	-	-	-	-	-
4500+ .....	-	-	-	-	-	-	-	-	-	-	-
Unknown .....	-	-	-	-	-	-	-	-	-	-	-

- Quantity is zero.

<sup>1</sup> Late neonatal deaths occur from day 7 through 27 after birth.

NOTE: Calculated gestation from reported date of last menses. If calculated gestation is unknown, the clinical estimate of gestation is used. Total includes reports with unknown birthweight and/or gestation.