

**TABLE 7-10. Postneonatal Deaths<sup>1</sup> by Weeks of Gestation and Weight  
Oregon Residents, Birth Cohort 2009**

Birthweight (In Grams)	Total	Weeks of Gestation									
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+	N.S.
<b>Total .....</b>	69	–	–	5	5	3	6	35	11	4	–
001-349 .....	–	–	–	–	–	–	–	–	–	–	–
350-499 .....	–	–	–	–	–	–	–	–	–	–	–
<500 .....	–	–	–	–	–	–	–	–	–	–	–
500-749 .....	5	–	–	4	1	–	–	–	–	–	–
750-999 .....	1	–	–	1	–	–	–	–	–	–	–
1000-1249 .....	2	–	–	–	2	–	–	–	–	–	–
1250-1499 .....	–	–	–	–	–	–	–	–	–	–	–
1500-1999 .....	5	–	–	–	2	1	1	1	–	–	–
2000-2499 .....	4	–	–	–	–	2	–	2	–	–	–
<2500 .....	17	–	–	5	5	3	1	3	–	–	–
2500+ .....	52	–	–	–	–	–	5	32	11	4	–
2500-2999 .....	17	–	–	–	–	–	4	11	–	2	–
3000-3499 .....	18	–	–	–	–	–	1	11	5	1	–
3500-3999 .....	13	–	–	–	–	–	–	7	6	–	–
4000-4499 .....	3	–	–	–	–	–	–	2	–	1	–
4500+ .....	1	–	–	–	–	–	–	1	–	–	–
Unknown .....	–	–	–	–	–	–	–	–	–	–	–

– Quantity is zero.

<sup>1</sup> Postneonatal deaths occur from day 28 through 364 after birth.

NOTE: Calculated gestation from reported date of last menses. If calculated gestation is unknown, the clinical estimate of gestation is used. Total includes reports with unknown birthweight and/or gestation.