

**TABLE 7-9. Late neonatal deaths<sup>1</sup> by weeks of gestation and weight, Oregon residents, birth cohort 2014**

Birthweight (in grams)	Total	Weeks of gestation <sup>2</sup>								
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+
Total <sup>3</sup> .....	23	–	1	11	–	3	1	6	1	–
001-349 .....	–	–	–	–	–	–	–	–	–	–
350-499 .....	–	–	–	–	–	–	–	–	–	–
<500 .....	–	–	–	–	–	–	–	–	–	–
500-749 .....	4	–	1	3	–	–	–	–	–	–
750-999 .....	7	–	–	7	–	–	–	–	–	–
1000-1249 .....	1	–	–	1	–	–	–	–	–	–
1250-1499 .....	–	–	–	–	–	–	–	–	–	–
1500-1999 .....	1	–	–	–	–	1	–	–	–	–
2000-2499 .....	2	–	–	–	–	2	–	–	–	–
<2500 .....	15	–	1	11	–	3	–	–	–	–
2500+ .....	8	–	–	–	–	–	1	6	1	–
2500-2999 .....	2	–	–	–	–	–	–	2	–	–
3000-3499 .....	1	–	–	–	–	–	–	–	1	–
3500-3999 .....	4	–	–	–	–	–	1	3	–	–
4000-4499 .....	1	–	–	–	–	–	–	1	–	–
4500+ .....	–	–	–	–	–	–	–	–	–	–

– Quantity is zero.

<sup>1</sup> Late neonatal deaths occur from day seven through 27 after birth.

<sup>2</sup> Clinical estimate of gestation. If clinical estimate is unknown, gestation is calculated from reported date of last menses.

<sup>3</sup> Total includes reports with unknown birthweight.