

**Table 1: Age-adjusted incidence rates of osteosarcoma (per million population), US and States, 2007-2011 combined (Source: CDC wonder)**

<u>Ages &lt;1-19 years</u>			<u>Ages &lt;1-24 years</u>		
State	Count	Rate	State	Count	Rate
<b>US</b>	<b>2,030</b>	<b>4.9</b>	<b>US</b>	<b>2,423</b>	<b>4.6</b>
Utah	36	8.1	Utah	40	7.2
Oregon	36	7.4	Iowa	34	6.6
Louisiana	42	6.8	Louisiana	49	6.3
Iowa	27	6.6	Oregon	37	6.1
Kansas	24	6.0	New Jersey	83	5.8
Florida	137	5.9	Florida	165	5.7
New Jersey	68	5.9	Idaho	16	5.7
California	281	5.4	Kansas	28	5.6
Texas	188	5.1	Michigan	89	5.1
Oklahoma	26	5.1	California	332	5.0
New York	127	5.0	New York	156	4.9
Michigan	70	5.0	Texas	222	4.9
North Carolina	62	4.9	Nevada	17	4.8
Massachusetts	42	4.9	Oklahoma	30	4.7
Indiana	43	4.7	Virginia	60	4.6
Illinois	84	4.7	New Mexico	16	4.5
Virginia	49	4.7	Illinois	100	4.5
Mississippi	20	4.6	Mississippi	24	4.5
Arkansas	17	4.4	North Carolina	70	4.5
Maryland	33	4.3	Indiana	48	4.3
Colorado	28	4.2	Massachusetts	45	4.2
Pennsylvania	69	4.2	Pennsylvania	86	4.1
Washington	37	4.2	Minnesota	37	4.1
Wisconsin	32	4.1	Ohio	79	4.1
South Carolina	25	4.1	Maryland	39	4.1
Tennessee	34	4.1	Tennessee	42	4.0
Alabama	26	4.0	Washington	44	3.9
Minnesota	28	3.9	Wisconsin	37	3.9
Missouri	31	3.9	Alabama	31	3.8
Ohio	60	3.9	Missouri	38	3.8
Arizona	32	3.6	Arizona	42	3.8
Georgia	48	3.5	South Carolina	29	3.8
Kentucky	20	3.5	Colorado	31	3.8
Connecticut	16	3.3	Arkansas	17	3.5
			Georgia	58	3.4
			Kentucky	24	3.3
			Connecticut	19	3.2

Source: United States Cancer Statistics: 1999 - 2011 Incidence, WONDER Online Database. United States Department of Health and Human Services, Centers for Disease Control and Prevention and National Cancer Institute; 2014. Accessed at <http://wonder.cdc.gov/cancer-v2011.html> on Oct 2, 2014 3:56:10 PM.

Caveats:

1. These states are missing data for the indicated year: Alabama 1999; Arkansas 1999-2000, 2002; District of Columbia 2001-2002; Maryland 2001; Mississippi 1999-2002; Nevada 1999, 2011; North Dakota 2001; South Dakota 1999-2000; Tennessee 1999-2003; Virginia 1999-2002.
2. Data are suppressed if fewer than 16 cases are reported in the specific category.
3. Data are from selected statewide and metropolitan area cancer registries that meet data quality criteria.
4. Information on primary site, behavior, and histology was coded according to the International Classification of Diseases for Oncology, Third Edition (ICD-O-3) and categorized according to the revised SEER recodes dated January 27, 2003, which define standard groupings of primary cancer sites.