

## Health screenings among adults by race and ethnicity, Oregon 2010-2011



Use this table to estimate the overall burden of disease within each race and ethnicity

	White, NL		African American, NL		Asian or Pacific Islander, NL		American Indian or Alaska Native, NL		Latino	
	Unadjusted (%)	Number of adults	Unadjusted (%)	Number of adults	Unadjusted (%)	Number of adults	Unadjusted (%)	Number of adults	Unadjusted (%)	Number of adults
Cholesterol checked within past five years	73.5	1,828,500	72.5	40,600	71.0	93,000	75.8	29,500	54.8	151,800
Current <sup>1</sup> on colorectal cancer screening (50-75 years old)	60.6	417,200	62.9	9,900	52.9	17,600	55.8	7,100	20.8	5,800
FOBT within one year (50-75 years old)	12.6	86,400	21.9	3,500	8.6 ^	3,000	8.2	1,000	4.5 ^	1,300
Mammogram within past two years (women 50-74 years old)	74.8	261,100	73.3	5,200	79.1	14,100	63.3	4,500	72.3	9,600
Pap test within past three years (women 21-65 years old with a cervix)	82.2	446,400	87.6	13,700	76.1	34,300	75.7	8,400	85.1	59,100

FOBT = Fecal occult blood test; NL = non-Latino

^ This number may be statistically unreliable and should be interpreted with caution.

<sup>1</sup> Current on colorectal cancer screening includes: having a fecal occult blood test (FOBT) in the past year; a colonoscopy within the past 10 years; or, a sigmoidoscopy within the past five years as well as an FOBT within the past three years.

## Health screenings among adults by race and ethnicity, Oregon 2010-2011

Use this table to compare burden of disease among race and ethnicity groups

	White, NL (%)	African American, NL (%)	Asian or Pacific Islander, NL (%)	American Indian or Alaska Native, NL (%)	Latino (%)
Cholesterol checked within past five years*	69.5	76.1	73.5	76.6	59.9
Current <sup>1</sup> on colorectal cancer screening (50-75 years old)	60.6	62.9	52.9	55.8	20.8
FOBT within one year (50-75 years old)	12.6	21.9	8.6 ^	8.2	4.5 ^
Mammogram within past two years (women 50-74 years old)	74.8	73.3	79.1	63.3	72.3
Pap test within past three years (women 21-65 years old with a cervix)*	82.4	86.3	74.9	77.3	76.5

FOBT = Fecal occult blood test; NA = not applicable or available; NL = non-Latino

^ This number may be statistically unreliable and should be interpreted with caution.

\* Age-adjusted to the 2000 standard population.

<sup>1</sup> Current on colorectal cancer screening includes: having a fecal occult blood test (FOBT) in the past year; a colonoscopy within the past 10 years; or, a sigmoidoscopy within the past five years as well as an FOBT within the past three years.

Source: Oregon Behavioral Risk Factors Surveillance System Race Oversample Dataset 2010-2011; age-adjusted to the 2000 standard population.

Note: For an explanation of using unadjusted versus age-adjusted estimates, please see <http://public.health.oregon.gov/DiseasesConditions/ChronicDisease/DataReports/Pages/TechnicalNotes.aspx>.

Suggested citation: Oregon Health Authority, Public Health Division, Health Promotion and Chronic Disease Prevention section. Health screenings among adults by race and ethnicity, Oregon 2010-2011. <http://public.health.oregon.gov/DiseasesConditions/ChronicDisease/DataReports/Pages/AdultData.aspx>.

Created August 11, 2014. Accessed [DOWNLOAD DATE].