

Oregon Tuberculosis Control Program Indicators

1. Reporting and Treatment for TB cases

Outcome	Year							
	2000	2001	2002	2003	2004	2005	2006	2007
Completion of therapy within one year [†]	81%	88%	87%	84%	90%	94%	93%	91%
HIV status reported	81%	76%	77%	77%	82%	93%	78%	94%
Totally directly observed therapy	55%	64%	68%	70%	61%	65%	67%	65%

2. Follow-up and Treatment for Contacts to Sputum-smear Positive Cases

Outcome	Year							
	2000	2001	2002	2003	2004	2005	2006*	2007*
Cases with no contacts	2%	11%	7%	6%	7%	6%	7%	7%
Contacts evaluated for TB	92%	84%	81%	88%	75%	71%	87%	80%
Active TB identified	2%	2%	1%	1%	0%	1%	2%	2%
NEW Latent TB identified [‡]	43%	39%	37%	32%	23%	19%	22%	24%
LTBI treatment started [‡]	71%	93%	92%	87%	80%	76%	86%	76%
LTBI treatment completed [‡]	Not Avail.	85%	73%	79%	58%	69%	84%	75%

Updated 10/2009

[†]Excludes patients with resistance to rifampin, patients who died during treatment, patients with meningeal TB, children under age of 15 (at report) with disseminated TB (defined as miliary and/or positive blood culture).

[‡] Excludes contacts with prior positive TST/QFT test and/or prior history of treatment for TB infection or disease.

*Oregon data from 2006 on include contact investigations to sputum smear positive AND/OR cavitory and laryngeal cases.