


Early Syphilis Cases by Year, Oregon, 2006–2015




Cumulative Counts, Proportions and Rates for 2006–2015

	(Count)	(%)	(per 100,000)
Age			
10 – 14 yr.	0	0	0
15 – 19 yr.	44	2.1	1.8
20 – 24 yr.	218	10.6	.6
25 – 34 yr.	316	15.4	.8
35 – 44 yr.	578	28.2	11.2
45 – 54 yr.	500	24.4	9.8
55 – 64 yr.	303	14.8	5.6
≥ 65 yr.	93	4.5	.2
Sex			
Male	1936	94.3	10.2
Female	116	5.7	.6
Race, ethnicity			
Non-hispanic white	1409	69.8	46.1
Non-hispanic black	118	5.8	166.4
Non-hispanic native American, Alaska native	18	.9	40.5
Non-hispanic Pacific Islander	8	.4	53.4
Non-hispanic Asian	44	2.2	26.8
Hispanic	340	16.8	68.4
Other	11	.5	n/a
Multiple	0	0	0
Refused, unknown	70	3.5	n/a
Risk			
Male, sex with males	1231	63.6*	n/a
Other Infections			n/a
HIV	843	41.1	


*Represents male cases that were asked and answered 'Yes' to question about ever having sex with men as a proportion of all male cases reported .

Early Syphilis Cases by Year, Baker, 2006–2015



The Oregon Health Division HIV Surveillance Program does not publish counts of syphilis cases by age, race, sex or transmission group if fewer than 10 cases have been reported during the above time interval, county population is less than 10,000, or the estimated county population of any race group or age group typically used to group cases in its reports is smaller than 50. If you are a public health official from this county and would like to request estimates of syphilis cases in your county by age and race and sex, please contact the Oregon Health Division STD Program

Early Syphilis Cases by Year, Benton, 2006–2015




Cumulative Counts, Proportions and Rates for 2006–2015

	(Count)	(%)	(per 100,000)
Age			
10 – 14 yr.	0	0	0
15 – 19 yr.	1	5	1.2
20 – 24 yr.	6	30	.7
25 – 34 yr.	5	25	.6
35 – 44 yr.	4	20	4.5
45 – 54 yr.	2	10	2.2
55 – 64 yr.	1	5	.9
≥ 65 yr.	1	5	.1
Sex			
Male	20	100	4.7
Female	0	0	0
Race, ethnicity			
Non-hispanic white	13	65	18.4
Non-hispanic black	1	5	114.9
Non-hispanic native American, Alaska native	0	0	0
Non-hispanic Pacific Islander	0	0	0
Non-hispanic Asian	0	0	0
Hispanic	5	25	82.4
Other	0	0	n/a
Multiple	0	0	0
Refused, unknown	1	5	n/a
Risk			
Male, sex with males	13	65*	n/a
Other Infections			n/a
HIV	2	10	

*Represents male cases that were asked and answered 'Yes' to question about ever having sex with men as a proportion of all male cases reported .

Early Syphilis Cases by Year, Clackamas, 2006–2015




Cumulative Counts, Proportions and Rates for 2006–2015

	(Count)	(%)	(per 100,000)
Age			
10 – 14 yr.	0	0	0
15 – 19 yr.	0	0	0
20 – 24 yr.	8	5.8	.2
25 – 34 yr.	18	13.1	.5
35 – 44 yr.	32	23.4	6.9
45 – 54 yr.	42	30.7	7.6
55 – 64 yr.	30	21.9	5
≥ 65 yr.	7	5.1	.2
Sex			
Male	127	92.7	6.8
Female	10	7.3	.5
Race, ethnicity			
Non-hispanic white	106	77.9	32.3
Non-hispanic black	5	3.7	147.8
Non-hispanic native American, Alaska native	1	.7	40.3
Non-hispanic Pacific Islander	0	0	0
Non-hispanic Asian	3	2.2	18.4
Hispanic	15	11	45.5
Other	1	.7	n/a
Multiple	0	0	0
Refused, unknown	5	3.7	n/a
Risk			
Male, sex with males	64	50.4*	n/a
Other Infections			n/a
HIV	56	40.6	


*Represents male cases that were asked and answered 'Yes' to question about ever having sex with men as a proportion of all male cases reported .

Early Syphilis Cases by Year, Clatsop, 2006–2015




The Oregon Health Division HIV Surveillance Program does not publish counts of syphilis cases by age, race, sex or transmission group if fewer than 10 cases have been reported during the above time interval, county population is less than 10,000, or the estimated county population of any race group or age group typically used to group cases in its reports is smaller than 50. If you are a public health official from this county and would like to request estimates of syphilis cases in your county by age and race and sex, please contact the Oregon Health Division STD Program

Early Syphilis Cases by Year, Columbia, 2006–2015




The Oregon Health Division HIV Surveillance Program does not publish counts of syphilis cases by age, race, sex or transmission group if fewer than 10 cases have been reported during the above time interval, county population is less than 10,000, or the estimated county population of any race group or age group typically used to group cases in its reports is smaller than 50. If you are a public health official from this county and would like to request estimates of syphilis cases in your county by age and race and sex, please contact the Oregon Health Division STD Program

Early Syphilis Cases by Year, Coos, 2006–2015




The Oregon Health Division HIV Surveillance Program does not publish counts of syphilis cases by age, race, sex or transmission group if fewer than 10 cases have been reported during the above time interval, county population is less than 10,000, or the estimated county population of any race group or age group typically used to group cases in its reports is smaller than 50. If you are a public health official from this county and would like to request estimates of syphilis cases in your county by age and race and sex, please contact the Oregon Health Division STD Program

Early Syphilis Cases by Year, Crook, 2006–2015




The Oregon Health Division HIV Surveillance Program does not publish counts of syphilis cases by age, race, sex or transmission group if fewer than 10 cases have been reported during the above time interval, county population is less than 10,000, or the estimated county population of any race group or age group typically used to group cases in its reports is smaller than 50. If you are a public health official from this county and would like to request estimates of syphilis cases in your county by age and race and sex, please contact the Oregon Health Division STD Program

Early Syphilis Cases by Year, Curry, 2006–2015



The Oregon Health Division HIV Surveillance Program does not publish counts of syphilis cases by age, race, sex or transmission group if fewer than 10 cases have been reported during the above time interval, county population is less than 10,000, or the estimated county population of any race group or age group typically used to group cases in its reports is smaller than 50. If you are a public health official from this county and would like to request estimates of syphilis cases in your county by age and race and sex, please contact the Oregon Health Division STD Program

Early Syphilis Cases by Year, Deschutes, 2006–2015




Cumulative Counts, Proportions and Rates for 2006–2015

	(Count)	(%)	(per 100,000)
Age			
10 – 14 yr.	0	0	0
15 – 19 yr.	2	8.7	2.1
20 – 24 yr.	8	34.8	.5
25 – 34 yr.	5	21.7	.3
35 – 44 yr.	4	17.4	1.9
45 – 54 yr.	4	17.4	1.8
55 – 64 yr.	0	0	0
≥ 65 yr.	0	0	0
Sex			
Male	18	78.3	2.3
Female	5	21.7	.6
Race, ethnicity			
Non-hispanic white	16	80	10.7
Non-hispanic black	0	0	0
Non-hispanic native American, Alaska native	0	0	0
Non-hispanic Pacific Islander	0	0	0
Non-hispanic Asian	0	0	0
Hispanic	3	15	22.6
Other	0	0	n/a
Multiple	0	0	0
Refused, unknown	1	5	n/a
Risk			
Male, sex with males	9	50*	n/a
Other Infections			n/a
HIV	2	8.7	

*Represents male cases that were asked and answered 'Yes' to question about ever having sex with men as a proportion of all male cases reported .

Early Syphilis Cases by Year, Douglas, 2006–2015




Cumulative Counts, Proportions and Rates for 2006–2015

	(Count)	(%)	(per 100,000)
Age			
10 – 14 yr.	0	0	0
15 – 19 yr.	3	10.7	4.5
20 – 24 yr.	10	35.7	.9
25 – 34 yr.	6	21.4	.6
35 – 44 yr.	5	17.9	4.6
45 – 54 yr.	4	14.3	3.1
55 – 64 yr.	0	0	0
≥ 65 yr.	0	0	0
Sex			
Male	16	57.1	3
Female	12	42.9	2.2
Race, ethnicity			
Non-hispanic white	26	96.3	27.4
Non-hispanic black	0	0	0
Non-hispanic native American, Alaska native	1	3.7	55.1
Non-hispanic Pacific Islander	0	0	0
Non-hispanic Asian	0	0	0
Hispanic	0	0	0
Other	0	0	n/a
Multiple	0	0	0
Refused, unknown	0	0	n/a
Risk			
Male, sex with males	5	31.3*	n/a
Other Infections			n/a
HIV	2	7.1	


*Represents male cases that were asked and answered 'Yes' to question about ever having sex with men as a proportion of all male cases reported .

Early Syphilis Cases by Year, Gilliam, 2006–2015




The Oregon Health Division HIV Surveillance Program does not publish counts of syphilis cases by age, race, sex or transmission group if fewer than 10 cases have been reported during the above time interval, county population is less than 10,0000, or the estimated county population of any race group or age group typically used to group cases in its reports is smaller than 50. If you are a public health official from this county and would like to request estimates of syphilis cases in your county by age and race and sex, please contact the Oregon Health Division STD Program

Early Syphilis Cases by Year, Grant, 2006–2015




The Oregon Health Division HIV Surveillance Program does not publish counts of syphilis cases by age, race, sex or transmission group if fewer than 10 cases have been reported during the above time interval, county population is less than 10,000, or the estimated county population of any race group or age group typically used to group cases in its reports is smaller than 50. If you are a public health official from this county and would like to request estimates of syphilis cases in your county by age and race and sex, please contact the Oregon Health Division STD Program

Early Syphilis Cases by Year, Harney, 2006–2015




The Oregon Health Division HIV Surveillance Program does not publish counts of syphilis cases by age, race, sex or transmission group if fewer than 10 cases have been reported during the above time interval, county population is less than 10,0000, or the estimated county population of any race group or age group typically used to group cases in its reports is smaller than 50. If you are a public health official from this county and would like to request estimates of syphilis cases in your county by age and race and sex, please contact the Oregon Health Division STD Program

Early Syphilis Cases by Year, Hood River, 2006–2015



The Oregon Health Division HIV Surveillance Program does not publish counts of syphilis cases by age, race, sex or transmission group if fewer than 10 cases have been reported during the above time interval, county population is less than 10,000, or the estimated county population of any race group or age group typically used to group cases in its reports is smaller than 50. If you are a public health official from this county and would like to request estimates of syphilis cases in your county by age and race and sex, please contact the Oregon Health Division STD Program

Early Syphilis Cases by Year, Jackson, 2006–2015




Cumulative Counts, Proportions and Rates for 2006–2015

	(Count)	(%)	(per 100,000)
Age			
10 – 14 yr.	0	0	0
15 – 19 yr.	0	0	0
20 – 24 yr.	7	21.9	.3
25 – 34 yr.	4	12.5	.2
35 – 44 yr.	10	31.3	4.3
45 – 54 yr.	4	12.5	1.6
55 – 64 yr.	5	15.6	1.6
≥ 65 yr.	2	6.3	.1
Sex			
Male	26	81.3	2.6
Female	6	18.8	.6
Race, ethnicity			
Non-hispanic white	24	75	13.9
Non-hispanic black	1	3.1	69.1
Non-hispanic native American, Alaska native	1	3.1	51.4
Non-hispanic Pacific Islander	0	0	0
Non-hispanic Asian	0	0	0
Hispanic	4	12.5	15.8
Other	0	0	n/a
Multiple	0	0	0
Refused, unknown	2	6.3	n/a
Risk			
Male, sex with males	8	30.8*	n/a
Other Infections			n/a
HIV	3	9.4	


*Represents male cases that were asked and answered 'Yes' to question about ever having sex with men as a proportion of all male cases reported .

Early Syphilis Cases by Year, Jefferson, 2006–2015



The Oregon Health Division HIV Surveillance Program does not publish counts of syphilis cases by age, race, sex or transmission group if fewer than 10 cases have been reported during the above time interval, county population is less than 10,0000, or the estimated county population of any race group or age group typically used to group cases in its reports is smaller than 50. If you are a public health official from this county and would like to request estimates of syphilis cases in your county by age and race and sex, please contact the Oregon Health Division STD Program

Early Syphilis Cases by Year, Josephine, 2006–2015




Cumulative Counts, Proportions and Rates for 2006–2015

	(Count)	(%)	(per 100,000)
Age			
10 – 14 yr.	0	0	0
15 – 19 yr.	0	0	0
20 – 24 yr.	1	7.7	.1
25 – 34 yr.	1	7.7	.1
35 – 44 yr.	5	38.5	6.2
45 – 54 yr.	1	7.7	1
55 – 64 yr.	2	15.4	1.6
≥ 65 yr.	3	23.1	.4
Sex			
Male	13	100	3.2
Female	0	0	0
Race, ethnicity			
Non-hispanic white	8	66.7	10.9
Non-hispanic black	0	0	0
Non-hispanic native American, Alaska native	0	0	0
Non-hispanic Pacific Islander	0	0	0
Non-hispanic Asian	0	0	0
Hispanic	1	8.3	17.1
Other	0	0	n/a
Multiple	0	0	0
Refused, unknown	3	25	n/a
Risk			
Male, sex with males	5	38.5*	n/a
Other Infections			n/a
HIV	2	15.4	


*Represents male cases that were asked and answered 'Yes' to question about ever having sex with men as a proportion of all male cases reported .

Early Syphilis Cases by Year, Klamath, 2006–2015



The Oregon Health Division HIV Surveillance Program does not publish counts of syphilis cases by age, race, sex or transmission group if fewer than 10 cases have been reported during the above time interval, county population is less than 10,000, or the estimated county population of any race group or age group typically used to group cases in its reports is smaller than 50. If you are a public health official from this county and would like to request estimates of syphilis cases in your county by age and race and sex, please contact the Oregon Health Division STD Program

Early Syphilis Cases by Year, Lake, 2006–2015



The Oregon Health Division HIV Surveillance Program does not publish counts of syphilis cases by age, race, sex or transmission group if fewer than 10 cases have been reported during the above time interval, county population is less than 10,0000, or the estimated county population of any race group or age group typically used to group cases in its reports is smaller than 50. If you are a public health official from this county and would like to request estimates of syphilis cases in your county by age and race and sex, please contact the Oregon Health Division STD Program

Early Syphilis Cases by Year, Lane, 2006–2015




Cumulative Counts, Proportions and Rates for 2006–2015

	(Count)	(%)	(per 100,000)
Age			
10 – 14 yr.	0	0	0
15 – 19 yr.	3	3.2	1.2
20 – 24 yr.	18	19.4	.5
25 – 34 yr.	22	23.7	.6
35 – 44 yr.	24	25.8	5.6
45 – 54 yr.	11	11.8	2.6
55 – 64 yr.	11	11.8	2.2
≥ 65 yr.	4	4.3	.1
Sex			
Male	79	84.9	4.6
Female	14	15.1	.8
Race, ethnicity			
Non-hispanic white	67	72	22.5
Non-hispanic black	2	2.2	59.9
Non-hispanic native American, Alaska native	3	3.2	81.5
Non-hispanic Pacific Islander	0	0	0
Non-hispanic Asian	6	6.5	60
Hispanic	11	11.8	37.1
Other	0	0	n/a
Multiple	0	0	0
Refused, unknown	4	4.3	n/a
Risk			
Male, sex with males	42	53.2*	n/a
Other Infections			n/a
HIV	19	20.4	

*Represents male cases that were asked and answered 'Yes' to question about ever having sex with men as a proportion of all male cases reported .

Early Syphilis Cases by Year, Lincoln, 2006–2015




Cumulative Counts, Proportions and Rates for 2006–2015

	(Count)	(%)	(per 100,000)
Age			
10 – 14 yr.	0	0	0
15 – 19 yr.	2	16.7	8.3
20 – 24 yr.	3	25	.7
25 – 34 yr.	1	8.3	.2
35 – 44 yr.	1	8.3	2.2
45 – 54 yr.	3	25	5.6
55 – 64 yr.	1	8.3	1.2
≥ 65 yr.	1	8.3	.2
Sex			
Male	6	50	2.7
Female	6	50	2.5
Race, ethnicity			
Non-hispanic white	10	90.9	25.9
Non-hispanic black	1	9.1	384.6
Non-hispanic native American, Alaska native	0	0	0
Non-hispanic Pacific Islander	0	0	0
Non-hispanic Asian	0	0	0
Hispanic	0	0	0
Other	0	0	n/a
Multiple	0	0	0
Refused, unknown	0	0	n/a
Risk			
Male, sex with males	3	50*	n/a
Other Infections			n/a
HIV	1	8.3	

*Represents male cases that were asked and answered 'Yes' to question about ever having sex with men as a proportion of all male cases reported .

Early Syphilis Cases by Year, Linn, 2006–2015




Cumulative Counts, Proportions and Rates for 2006–2015

	(Count)	(%)	(per 100,000)
Age			
10 – 14 yr.	0	0	0
15 – 19 yr.	3	12	4
20 – 24 yr.	7	28	.6
25 – 34 yr.	6	24	.5
35 – 44 yr.	4	16	2.8
45 – 54 yr.	1	4	.7
55 – 64 yr.	3	12	1.8
≥ 65 yr.	1	4	.1
Sex			
Male	24	96	4.2
Female	1	4	.2
Race, ethnicity			
Non-hispanic white	16	64	15.5
Non-hispanic black	2	8	348.4
Non-hispanic native American, Alaska native	1	4	75.2
Non-hispanic Pacific Islander	0	0	0
Non-hispanic Asian	0	0	0
Hispanic	6	24	60.2
Other	0	0	n/a
Multiple	0	0	0
Refused, unknown	0	0	n/a
Risk			
Male, sex with males	19	79.2*	n/a
Other Infections			n/a
HIV	4	16	

*Represents male cases that were asked and answered 'Yes' to question about ever having sex with men as a proportion of all male cases reported .

Early Syphilis Cases by Year, Malheur, 2006–2015



The Oregon Health Division HIV Surveillance Program does not publish counts of syphilis cases by age, race, sex or transmission group if fewer than 10 cases have been reported during the above time interval, county population is less than 10,000, or the estimated county population of any race group or age group typically used to group cases in its reports is smaller than 50. If you are a public health official from this county and would like to request estimates of syphilis cases in your county by age and race and sex, please contact the Oregon Health Division STD Program

Early Syphilis Cases by Year, Marion, 2006–2015




Cumulative Counts, Proportions and Rates for 2006–2015

	(Count)	(%)	(per 100,000)
Age			
10 – 14 yr.	0	0	0
15 – 19 yr.	4	3.2	1.7
20 – 24 yr.	22	17.6	.7
25 – 34 yr.	15	12	.5
35 – 44 yr.	37	29.6	9
45 – 54 yr.	28	22.4	7
55 – 64 yr.	16	12.8	3.9
≥ 65 yr.	3	2.4	.1
Sex			
Male	113	90.4	7.2
Female	12	9.6	.8
Race, ethnicity			
Non-hispanic white	68	55.7	31.2
Non-hispanic black	6	4.9	178.1
Non-hispanic native American, Alaska native	1	.8	29.4
Non-hispanic Pacific Islander	4	3.3	145
Non-hispanic Asian	2	1.6	29.9
Hispanic	39	32	46.6
Other	1	.8	n/a
Multiple	0	0	0
Refused, unknown	1	.8	n/a
Risk			
Male, sex with males	79	69.9*	n/a
Other Infections			n/a
HIV	38	30.4	


*Represents male cases that were asked and answered 'Yes' to question about ever having sex with men as a proportion of all male cases reported .

Early Syphilis Cases by Year, Morrow, 2006–2015



The Oregon Health Division HIV Surveillance Program does not publish counts of syphilis cases by age, race, sex or transmission group if fewer than 10 cases have been reported during the above time interval, county population is less than 10,0000, or the estimated county population of any race group or age group typically used to group cases in its reports is smaller than 50. If you are a public health official from this county and would like to request estimates of syphilis cases in your county by age and race and sex, please contact the Oregon Health Division STD Program

Early Syphilis Cases by Year, Multnomah, 2006–2015




Cumulative Counts, Proportions and Rates for 2006–2015

	(Count)	(%)	(per 100,000)
Age			
10 – 14 yr.	0	0	0
15 – 19 yr.	12	1	3
20 – 24 yr.	82	6.9	1.1
25 – 34 yr.	173	14.6	2.3
35 – 44 yr.	345	29.1	26.9
45 – 54 yr.	319	26.9	30.4
55 – 64 yr.	197	16.6	20.1
≥ 65 yr.	57	4.8	.8
Sex			
Male	1158	97.7	31.7
Female	27	2.3	.7
Race, ethnicity			
Non-hispanic white	847	72.8	152.9
Non-hispanic black	85	7.3	206.6
Non-hispanic native American, Alaska native	8	.7	137.3
Non-hispanic Pacific Islander	3	.3	63.5
Non-hispanic Asian	23	2	41.8
Hispanic	164	14.1	188.5
Other	7	.6	n/a
Multiple	0	0	0
Refused, unknown	26	2.2	n/a
Risk			
Male, sex with males	834	72*	n/a
Other Infections			n/a
HIV	574	48.6	

*Represents male cases that were asked and answered 'Yes' to question about ever having sex with men as a proportion of all male cases reported .

Early Syphilis Cases by Year, Polk, 2006–2015



Cumulative Counts, Proportions and Rates for 2006–2015

	(Count)	(%)	(per 100,000)
Age			
10 – 14 yr.	0	0	0
15 – 19 yr.	0	0	0
20 – 24 yr.	3	21.4	.4
25 – 34 yr.	2	14.3	.3
35 – 44 yr.	3	21.4	3.5
45 – 54 yr.	5	35.7	5.5
55 – 64 yr.	1	7.1	1
≥ 65 yr.	0	0	0
Sex			
Male	13	92.9	3.6
Female	1	7.1	.3
Race, ethnicity			
Non-hispanic white	10	76.9	16.1
Non-hispanic black	0	0	0
Non-hispanic native American, Alaska native	0	0	0
Non-hispanic Pacific Islander	0	0	0
Non-hispanic Asian	0	0	0
Hispanic	2	15.4	19.8
Other	0	0	n/a
Multiple	0	0	0
Refused, unknown	1	7.7	n/a
Risk			
Male, sex with males	6	46.2*	n/a
Other Infections			n/a
HIV	4	28.6	


*Represents male cases that were asked and answered 'Yes' to question about ever having sex with men as a proportion of all male cases reported .

Early Syphilis Cases by Year, Sherman, 2006–2015




The Oregon Health Division HIV Surveillance Program does not publish counts of syphilis cases by age, race, sex or transmission group if fewer than 10 cases have been reported during the above time interval, county population is less than 10,0000, or the estimated county population of any race group or age group typically used to group cases in its reports is smaller than 50. If you are a public health official from this county and would like to request estimates of syphilis cases in your county by age and race and sex, please contact the Oregon Health Division STD Program

Early Syphilis Cases by Year, Tillamook, 2006–2015




The Oregon Health Division HIV Surveillance Program does not publish counts of syphilis cases by age, race, sex or transmission group if fewer than 10 cases have been reported during the above time interval, county population is less than 10,000, or the estimated county population of any race group or age group typically used to group cases in its reports is smaller than 50. If you are a public health official from this county and would like to request estimates of syphilis cases in your county by age and race and sex, please contact the Oregon Health Division STD Program

Early Syphilis Cases by Year, Umatilla, 2006–2015




The Oregon Health Division HIV Surveillance Program does not publish counts of syphilis cases by age, race, sex or transmission group if fewer than 10 cases have been reported during the above time interval, county population is less than 10,000, or the estimated county population of any race group or age group typically used to group cases in its reports is smaller than 50. If you are a public health official from this county and would like to request estimates of syphilis cases in your county by age and race and sex, please contact the Oregon Health Division STD Program

Early Syphilis Cases by Year, Union, 2006–2015




The Oregon Health Division HIV Surveillance Program does not publish counts of syphilis cases by age, race, sex or transmission group if fewer than 10 cases have been reported during the above time interval, county population is less than 10,0000, or the estimated county population of any race group or age group typically used to group cases in its reports is smaller than 50. If you are a public health official from this county and would like to request estimates of syphilis cases in your county by age and race and sex, please contact the Oregon Health Division STD Program

Early Syphilis Cases by Year, Wallowa, 2006–2015




The Oregon Health Division HIV Surveillance Program does not publish counts of syphilis cases by age, race, sex or transmission group if fewer than 10 cases have been reported during the above time interval, county population is less than 10,0000, or the estimated county population of any race group or age group typically used to group cases in its reports is smaller than 50. If you are a public health official from this county and would like to request estimates of syphilis cases in your county by age and race and sex, please contact the Oregon Health Division STD Program

Early Syphilis Cases by Year, Wasco, 2006–2015



The Oregon Health Division HIV Surveillance Program does not publish counts of syphilis cases by age, race, sex or transmission group if fewer than 10 cases have been reported during the above time interval, county population is less than 10,000, or the estimated county population of any race group or age group typically used to group cases in its reports is smaller than 50. If you are a public health official from this county and would like to request estimates of syphilis cases in your county by age and race and sex, please contact the Oregon Health Division STD Program

Early Syphilis Cases by Year, Washington, 2006–2015




Cumulative Counts, Proportions and Rates for 2006–2015

	(Count)	(%)	(per 100,000)
Age			
10 – 14 yr.	0	0	0
15 – 19 yr.	10	3.6	2.9
20 – 24 yr.	33	11.7	.6
25 – 34 yr.	47	16.7	.9
35 – 44 yr.	88	31.3	10.4
45 – 54 yr.	62	22.1	7.9
55 – 64 yr.	29	10.3	4.2
≥ 65 yr.	12	4.3	.2
Sex			
Male	266	94.7	10.1
Female	15	5.3	.6
Race, ethnicity			
Non-hispanic white	159	56.8	41.6
Non-hispanic black	13	4.6	124.3
Non-hispanic native American, Alaska native	1	.4	35.7
Non-hispanic Pacific Islander	1	.4	39.4
Non-hispanic Asian	8	2.9	14.7
Hispanic	75	26.8	81.8
Other	2	.7	n/a
Multiple	0	0	0
Refused, unknown	21	7.5	n/a
Risk			
Male, sex with males	120	45.1*	n/a
Other Infections			n/a
HIV	121	43.4	


*Represents male cases that were asked and answered 'Yes' to question about ever having sex with men as a proportion of all male cases reported .

Early Syphilis Cases by Year, Wheeler, 2006–2015



The Oregon Health Division HIV Surveillance Program does not publish counts of syphilis cases by age, race, sex or transmission group if fewer than 10 cases have been reported during the above time interval, county population is less than 10,0000, or the estimated county population of any race group or age group typically used to group cases in its reports is smaller than 50. If you are a public health official from this county and would like to request estimates of syphilis cases in your county by age and race and sex, please contact the Oregon Health Division STD Program

Early Syphilis Cases by Year, Yamhill, 2006–2015



Cumulative Counts, Proportions and Rates for 2006–2015

	(Count)	(%)	(per 100,000)
Age			
10 – 14 yr.	0	0	0
15 – 19 yr.	0	0	0
20 – 24 yr.	3	15	.3
25 – 34 yr.	5	25	.5
35 – 44 yr.	6	30	4.8
45 – 54 yr.	4	20	3.1
55 – 64 yr.	1	5	.8
≥ 65 yr.	1	5	.1
Sex			
Male	16	80	3.2
Female	4	20	.8
Race, ethnicity			
Non-hispanic white	11	55	13.9
Non-hispanic black	0	0	0
Non-hispanic native American, Alaska native	0	0	0
Non-hispanic Pacific Islander	0	0	0
Non-hispanic Asian	0	0	0
Hispanic	6	30	37.5
Other	0	0	n/a
Multiple	0	0	0
Refused, unknown	3	15	n/a
Risk			
Male, sex with males	5	31.3*	n/a
Other Infections			n/a
HIV	3	15	

*Represents male cases that were asked and answered 'Yes' to question about ever having sex with men as a proportion of all male cases reported .