


## Binge drinking in the past 30 days among adults in regions covered by Coordinated Care Organizations, 2010–2013

Gray lines represent confidence intervals



Coordinated Care Organization <i>(Data represent entire population of CCO regions - not just enrolled population)</i>	Percent of adults (age-adjusted)	Number of respondents (N)
AllCare Health Plan	16.3%	1,990
Cascade Health Alliance	14.0%	409
Columbia Pacific	17.2%	746
Eastern Oregon	18.0%	1,213
FamilyCare	18.0%	8,515
Health Share of Oregon	18.0%	8,270
InterCommunity Health Network	20.0%	1,689
Jackson CareConnect	16.9%	1,231
PacificSource C.S. - Central OR	18.0%	1,155
PacificSource C.S. - Columbia Gorge	16.0%	112
PrimaryHealth of Josephine County	14.8%	739
Trillium Community Health Plan	19.1%	2,330
Umpqua Health Alliance	13.5%	709
Western Oregon Advanced Health	17.4%	663
Willamette Valley Community Health	16.5%	2,025
Yamhill County Care Organization	13.5%	1,002
<i>Oregon State</i>	<i>17.7%</i>	<i>21,672</i>

## About the Data

Data Source: Oregon Behavioral Risk Factor Surveillance System (BRFSS)

### Data Notes:

Adults = age 18+

Number of respondents (N) is unweighted; percentages are weighted

Age-Adjusted to standard U.S. 2000 population - 3 groups (18-34, 35-54, 55+)

Question: Binge drinking is defined as males who had five or more alcoholic drinks and females who had four or more alcoholic drinks on one or more occasions in the past 30 days.

Date: September 07, 2016