

Maternal and Child Health


Infant breastfeeding

Breast milk is the most complete form of nutrition for infants; its benefits for infants’ health, growth, immunity, and development are well documented. The American Academy of Pediatrics recommends breastfeeding for reduced risk of infection in infants and for the prevention of childhood obesity. Breastfeeding enhances bonding between infant and mother.

Breastfeeding rates in Oregon are higher than in the U.S. as a whole (Figure 1). In 2013, 80.2% of Oregon mothers breastfed their infants at 8 weeks after delivery compared to 64.1% nationally. Oregon also has the second highest rate of breastfeeding at 6 months postpartum of any state: 68.2% in Oregon compared to 51.8% in the U.S. in 2013¹.

FIGURE 1

Breastfeeding 8 weeks after delivery by year, Oregon and U.S.




Source: CDC PRAMStat

¹ [Breastfeeding Among U.S. Children Born 2002-2013, CDC National Immunization Survey](#)

Infant breastfeeding overall in Oregon is high, but mothers who are <25 years old or low-income have lower rates (Figures 2 and 3). In the weeks following birth, mothers often reduce or stop breastfeeding for a variety of reasons. With active support, these barriers can be resolved so mothers can achieve their personal breastfeeding goals and meet breastfeeding recommendations.

FIGURE 2


Breastfeeding 8 weeks after delivery by income, Oregon, 2013



Source: Oregon Pregnancy Risk Assessment Monitoring System (PRAMS)

FIGURE 3

Breastfeeding 8 weeks after delivery by maternal age, Oregon, 2013



Source: Oregon Pregnancy Risk Assessment Monitoring System (PRAMS)

In Oregon, women who identify as Asian had the highest percentage of breastfeeding their babies at 8 weeks of age (87.5%), and Pacific Islander women the lowest (68.7%). Oregon conducts an ongoing statewide survey of mothers of newborns and of toddlers to better understand mothers' knowledge and experiences of breastfeeding and to determine where breastfeeding promotion efforts can best be targeted. Oregon WIC offers additional support through Breastfeeding Peer Counseling Programs and provides breast pumps when they are needed for breastfeeding support or for mothers returning to work or school.

Additional Resources: [Breastfeeding](#)

About the Data: Data is from the CDC PRAMStat System and the Oregon Pregnancy Risk Assessment Monitoring System (PRAMS) by year of birth. Unknowns are excluded from the analysis. Percentage of infants breastfed includes any breastfeeding of the infant at 8 weeks postpartum.

For More Information Contact: Claudia Bingham, claudia.w.bingham@state.or.us or Alfredo Sandoval, alfredo.p.sandoval@state.or.us

Date Updated: August 9, 2016

[Oregon State Health Profile](#)

OHA 9153-D (Rev) 09/13: This document can be provided upon request in an alternate format for individuals with disabilities or in a language other than English for people with limited English skills. To request this publication in another format or language, contact the Publications and Design Section at 503-378-3486, 711 for TTY, or email dhs-oha.publicationrequest@state.or.us.